



Faculty of Resource Science and Technology

**DEVELOPMENT OF SEROLOGICAL ASSAY FOR DETECTION OF
CHIKUNGUNYA VIRUS INFECTION**

**OOI KA BENG
(48675)**

**Bachelor of Science with Honours
(Resource Biotechnology)
2018**

Development of serological assay for detection of Chikungunya virus infection

Ooi Ka Beng

48675

A thesis submitted in fulfilment of the requirement for the degree of
Bachelor of Science with Honours

Supervisor: Dr. Magdline Sia Henry Sum
(Institute of Health and Community Medicine)

Co-supervisor: Madam Anna Andrew
(Faculty of Medicine and Health Sciences)

Resource Biotechnology

Faculty of Resource Science and Technology
UNIVERSITY MALAYSIA SARAWAK
2018

Grade: _____

Please tick (✓)
Final Year Project Report
Masters
PhD

✓

DECLARATION OF ORIGINAL WORK


This declaration is made on the4..... day of June.....2018.

Student's Declaration:

I, Ooi Ka Beng, 48675, Faculty of Resource Science and Technology hereby declare that the work entitled 'Development of Serological Assay for Detection of Chikungunya Virus Infection' is my original work. I have not copied from any other students' work or from any other sources except where due reference or acknowledgement is made explicitly in the text, nor has any part been written for me by another person.

04/06/2018


Date submitted



Ooi Ka Beng (48675)

Supervisor's Declaration:

I, Dr. Magdline Sia Henry Sum hereby certifies that the work entitled 'Development Of Serological Assay For Detection Of Chikungunya Virus Infection' was prepared by the above named student, and was submitted to the "FACULTY" as a * partial/full fulfillment for the conferment of Bachelor Of Science With Honours In Resource Biotechnology, and the aforementioned work, to the best of my knowledge, is the said student's work.

Received for examination by: 

(Dr. Magdline Sia Henry Sum)

Date: 04/06/2018

I declare that Project/Thesis is classified as (Please tick (√)):

- ☐ **CONFIDENTIAL** (Contains confidential information under the Official Secret Act 1972)*
☐ **RESTRICTED** (Contains restricted information as specified by the organization where research was done)*
☒ **OPEN ACCESS**

I declare this Project/Thesis is to be submitted to the Centre for Academic Information Services and uploaded into UNIMAS Institutional Repository (UNIMAS IR) (Please tick (√)):


- ☐ **YES**
☐ **NO**

Validation of Project/Thesis

I therefore duly affirm with free consent and willingly declare that this said Project/Thesis shall be placed officially in the Centre for Academic Information Services with the abiding interest and rights as follows:

- This Project/Thesis is the sole legal property of University Malaysia Sarawak (UNIMAS).
- The Centre for Academic Information Services has the lawful right to make copies for the purpose of academic and research only and not for other purpose.
- The Centre for Academic Information Services has the lawful right to digitalize the content for the Local Content Database.
- The Centre for Academic Information Services has the lawful right to make copies of the Project/Thesis for academic exchange between Higher Learning Institute.
- No dispute or any claim shall arise from the student itself neither third party on this Project/Thesis once it becomes the sole property of UNIMAS.
- This Project/Thesis or any material, data and information related to it shall not be distributed, published or disclosed to any party by the student except with UNIMAS permission.

Student signature



(9 June 2018)

Supervisor signature:



(9 June 2018)

Current Address:

Fakulti Sains dan Teknologi Sumber,
Universiti Malaysia Sarawak,
94300 Kota Samarahan,
Sarawak.

Notes: * If the Project/Thesis is **CONFIDENTIAL** or **RESTRICTED**, please attach together as annexure a letter from the organization with the period and reasons of confidentiality and restriction.

[The instrument was prepared by The Centre for Academic Information Services]

ACKNOWLEDGEMENT

Foremost, I would like to thank all the people who contributed to the work described in this thesis. I would like to express my sincere gratitude to my supervisor Dr. Magdline Sia Henry Sum for the constant guidance, advice, support of my study and research. Her guidance helped me in all the time of research and writing of this thesis. Besides my supervisor, I would like to thank my co-supervisor Madam Anna Andrew for the guidance, advice, patience, and imparting her knowledge and expertise in this study.

Last but not least, I would like to thank my beloved parents, Ooi Eng San and Tan Siew Mooi for supporting me spiritually, financially and words of encouragement throughout this project.

Furthermore, I would like to take this opportunity to thank the staffs of the Institute of Health and Community Medicine (IHCM), UNIMAS, post-graduate students and my lab mate Low Yong Chun, for their kind help, guidance and working together with me in the lab.

Development of serological assay for detection of Chikungunya virus infection

Ooi Ka Beng

Resource Biotechnology Programme
Faculty of Resource Science and Technology
University Malaysia Sarawak

ABSTRACT

Chikungunya virus (CHIKV) is a mosquito-borne virus that infects human. The infection caused by CHIKV and dengue virus (DENV) share some common signs and symptoms which includes high fever, chills, rashes, joint and muscle pain. However, the prognosis, patient care, and persistent symptoms are different. The accurate identification of CHIKV is important during CHIKV outbreak, since dengue infection is endemic in Malaysia. To date, the diagnosis of CHIKV could only be done at Institute of Medical Research (IMR) and National Public Health Laboratory (NPHL) which are located at Peninsular Malaysia. Hence, the aim of this study is to develop a serological assay for the detection of CHIKV infection by the use of a recombinant antigen. The recombinant CHIKV E2 protein are expressed in *Escherichia coli* (*E. coli*) using pET SUMO expression system. The expressed protein was evaluated by SDS-PAGE and western blot. Then the E2 recombinant protein was purified through ProBond Purification System (Novex, CA, USA) and dialysed gradually in PBS. Then, ELISA are carried out by using the E2 recombinant protein against various serum and the sensitivity and specificity are 60.7% and 90% respectively.

Keywords: CHIKV, recombinant CHIKV E2 protein, ELISA, sensitivity, specificity.

ABSTRAK

*Virus Chikungunya (CHIKV) adalah penyakit yang menjangkiti manusia melalui nyamuk. Jangkitan yang disebabkan oleh CHIKV dan denggi virus (DENV) berkongsi beberapa tanda biasa dan gejala termasuk demam, menggigil, ruam, sakit sendi dan otot. Walau bagaimanapun, prognosis, penjagaan pesakit, dan symptom berterusan adalah berbeza. Pengenalan tepat CHIKV adalah penting semasa wabak CHIKV, memandangkan jangkitan denggi adalah endemic di Malaysia,. Setakat ini, diagnosis jangkitan CHIKV hanya boleh dijalankan di Institut Penyelidikan Perubatan (IMR) dan Makmal Kesihatan Awam Kebangsaan (NPHL) yang terletak di Semenanjung Malaysia. Oleh itu, tujuan kajian ini adalah untuk membangunkan assay serologi untuk mengesan jangkitan CHIKV dengan menggunakan antigen rekombinan. Rekombinan protein CHIKV E2 telah diekspres di dalam *Escherichia coli* (*E.coli*) dengan menggunakan sistem ekspresi pET SUMO. Protein yang diekspres telah dinilai dengan menggunakan SDS-PAGE dan Western blot. E2 rekombinan protein telah dipurifikasi dengan menggunakan ProBond Purification System (Novex, CA, USA) dan dialysed dengan beransur-ansur dalam PBS. Kemudian, ELISA dijalankan dengan menggunakan protein rekombinan E2 terhadap pelbagai serum dan sensitivity dan spesifisiti adalah 60.7% dan 90% masing-masing.*

Kata kunci: CHIKV, rekombinan CHIKV E2 protein, ELISA, sensitivity, spesifisiti

TABLE OF CONTENTS

DECLARATIONI

ACKNOWLEDGEMENT III

ABSTRACTIV

ABSTRAKIV

TABLE OF CONTENTS V

List of Figures VII

List of Tables..... VIII

List of AbbreviationsIX

1.0 INTRODUCTION 1

2.0 LITERATURE REVIEW 3

 2.1 Chikungunya Virus3

 2.2 Genome of CHIKV 3

 2.3 E2 protein of CHIKV6

 2.4 Epidemiology of CHIKV6

 2.5 Clinical symptoms of CHIKV infection8

 2.6 Diagnosis of CHIKV infection8

 2.7 Treatment of CHIKV9

 2.8 Serological test..... 10

3.0 MATERIALS AND METHODS..... 11

 3.1 LB medium & agar plate preparation 11

 3.2 Bacteria culture growth..... 11

 3.3 Colony polymerase chain reaction (PCR)..... 11

 3.4 Selection of colony to culture overnight..... 13

 3.5 Preparation of glycerol stock for positive clone 13

 3.6 Plasmid DNA preparations 13

 3.7 Plasmid polymerase chain reaction (PCR)..... 13

 3.8 Transformation into BL21(DE3) One Shot Cells 14

3.9 Preparation of glycerol stock for transformed cell..... 14

3.10 Small scale protein expression..... 15

3.11 SDS-PAGE and Western blot 16

3.12 Protein Purification 17

3.13 Measurement of protein concentration 19

3.14 Gradient Dialysis 20

3.15 Enzyme-linked Immunosorbent Assay (ELISA)..... 20

4.0 RESULTS 21

4.1 Growing of the culture 21

4.2 Colony Polymerase Chain Reaction (PCR) 21

4.3 Transformation of E2 clone into BL21(DE3) One Shot Cells..... 23

4.4 Small scale protein expression..... 25

4.5 Protein purification 28

4.6 Protein concentration 30

4.7 Optimization of E2 protein concentration used 31

4.8 Enzyme-linked Immunosorbent Assay (ELISA) 33

4.9 Data analysis 36

5.0 DISCUSSION..... 39

6.0 CONCLUSION AND RECOMMENDATIONS..... 41

REFERENCES 42

APPENDICES..... 47

List of Figures

Figure 2.1 Genome of CHIKV	4
Figure 2.2 Phylogenetic tree of CHIKV	5
Figure 2.3 Worldwide distribution of CHIKV	7
Figure 4.1 Colony PCR-1	22
Figure 4.2 Colony PCR-2	22
Figure 4.3 Culture plate of E2 (COL3).....	23
Figure 4.4 Culture plate of E2 (COL5).....	23
Figure 4.5 Culture plate of vector pET SUMO	23
Figure 4.6 Analysis of E2 (COL3) & pET SUMO at different conditions	26
Figure 4.7 Analysis of E2 (COL5) & pET SUMO at different conditions.	27
Figure 4.8 Elution of purified E2 recombinant proteins.....	29
Figure 4.9 Elution of purified pET SUMO proteins.....	29
Figure 4.10 ELISA checkerboard.....	32
Figure 4.11 Graph of OD ₄₉₂ against antigen concentration	32
Figure 4.12 Template of ELISA plate probed with different sera.....	33
Figure 4.13 ELISA plates probed with different sera.....	34

List of Tables

Table 3.1 Primers for colony PCR.....	12
Table 3.2 Condition for colony PCR reaction	12
Table 4.1 Concentration of purified E2 recombinant protein with 8 ml of cell lysate	30
Table 4.2 Concentration of purified E2 recombinant protein with 16 ml of cell lysate	30
Table 4.3 Concentration of pooled dialyzed E2 recombinant protein	30
Table 4.4 Reading of ELISA checkerboard at OD ₄₉₂	31
Table 4.5 Reading of ELISA at OD ₄₉₂	35
Table 4.6 Positive control & negative control data tabulation	36
Table 4.7 Summary of the average of ELISA-1 and ELISA-2	37
Table 4.8 Sensitivity and specificity of serological assay	38

List of Abbreviations

ARB	Arbidol
BSA	Bovine serum albumin
CBB	Coomassie brilliant blue
CHIK	Chikungunya
CHIKV	Chikungunya virus
Colony PCR	Colony polymerase chain reaction
DNA	Deoxyribose nucleic acid
ELISA	Enzyme linked immunosorbent assay
HPC	Hepatitis C virus
IgG	Immunoglobulin G
IgM	Immunoglobulin M
IHCM	Institute of health and community medicine
IMR	Institute for Medical Research
IPTG	Isopropyl β-D-thiogalactoside
LB	Luria Bertani
NaCl	Sodium chloride
Nickel-HRP	Nickel with horseradish peroxidase
NPHL	National Public Health Laboratory

NSAIDs	Non-steroidal anti-inflammatory drugs
°C	Celcius
ORF	Open reading frame
PBS	Phosphate buffered saline
RNA	Ribose nucleic acid
R.O water	Reverse osmosis water
Rpm	Revolution per minute
RSV	Respiratory syncytial virus
RT-PCR	Real-time polymerase chain reaction
SDS-PAGE	Sodium dodecyl sulphate polyacrylamide gel electrophoresis
S.O.C	Super optimal broth with catabolite repression
SFV	Semliki forest virus
TAE	Tris acetate EDTA
UHQ water	Ultra high-quality water

1.0 INTRODUCTION

Chikungunya virus (CHIKV) is an arthropod-borne virus with major outbreaks in Africa, India, and South-East Asia (Power and Logue, 2007). CHIKV are transmitted by the same species of mosquitoes *Aedes aegypti* and *Aedes albopictus* (Thiboutot *et al.*, 2010). The major clinical symptom of CHIKV infection is febrile illness, which is clinically similar to symptoms of dengue virus infection (Karabatsos, 1975).

In Malaysia, molecular technique or immunofluorescence assay (IFA) are used for the detection of CHIKV infection. The molecular technique is used to detect the viral genome in the infected patient while IFA technique is used to detect the presence of antibodies against a specific antigen in serum. However, in Sarawak the validation of CHIKV infection can only be carried out through Institute for Medical Research (IMR) and National Public Health Laboratory (NPHL) at Peninsular Malaysia. Therefore, the identification of the infection will be delayed if an outbreak occurs because unlike dengue virus infection, CHIKV infection is rarely fatal and usually does not require close clinical supervision.

Due to the similar clinical symptoms of CHIKV infection with other virus infection such as dengue virus (DENV), CHIKV infection is often misdiagnosed (Morrison, 2014). Therefore, the ability to differentiate CHIKV infection from dengue virus infection would be important to have a proper control measures, particularly in areas where dengue virus infection is endemic or epidemic (Cho *et al.*, 2008). Hence, serological assay for the detection of CHIKV infection should be available in Sarawak. Thus, the aim of this study is to develop a serological assay for the detection of CHIKV infection.

Therefore, the objectives of this study are:

1. To express and purify the recombinant CHIKV E2 protein in *Escherichia coli* expression system.
2. To determine the specificity and sensitivity of serological assay developed for detection of CHIKV infection.

2.0 LITERATURE REVIEW

2.1 Chikungunya Virus

Chikungunya virus (CHIKV) belongs to the family *Togaviridae* and genus *Alphavirus*. The infection of CHIKV causes chikungunya fever (CHIK fever) which is a febrile illness linked with severe arthralgia and rash (McGill, 1995). The term 'Chikungunya' is derived from Makonde word meaning 'that which bends up' that refers to the contorted posture of the infected patients suffering from severe joint pain (Mavalankar *et al.*, 2008). CHIKV was first documented in 1952 during an epidemic outbreak in Makonde, Tanzania (Robinson, 1955). The transmission of CHIKV to humans are mostly through the bite of an infected *Aedes* mosquito. The transmission cycles of CHIKV includes urban cycle which is man – mosquito – man and sylvatic cycle animal – mosquito – man (Apandi *et al.*, 2009). The main vector that involves in urban cycles are usually *Aedes aegypti* and *Aedes albopictus* and *Aedes africanus*, *Aedes luteocephalus*, *Aedes furcifer* and *Aedes taylori* are the major vector involves in sylvatic cycle (Thiboutot *et al.*, 2010).

2.2 Genome of CHIKV

The genome of CHIKV (Figure 2.1) is a linear, single stranded, positive-sense RNA virus with approximately 11.8 kb and is capped in 5' and polyA tail in 3' end. The genome includes two open reading frames (ORFs) and encodes 4 non-structural proteins (nsP1, nsP2, nsP3 and nsP4) and five structural proteins (Capsid, E3, E2, 6k and E1) (Strauss *et al.*, 1991). The structural protein E1 and E2 associate as trimers of heterodimers (E2 – E1) as spikes on viral surface (Voss *et al.*, 2010) and the protein cover the viral surface in the form of membrane-anchored types (Cho *et al.*, 2008). The viral spikes help the attachment of the virus to cell surfaces and viral entry into the cells. The phylogenetic trees of CHIKV (Figure 2.2) shows that there are three

different genotypes based on geographical origins which are ECSA (East/Central/South African), West Africa, and Asian (Presti *et al.*, 2014). According to Schuffenecker *et al.* (2006), the single point mutation of alanine to valine at position 226 in E1 envelope glycoprotein (E1 – A226V1) during the Indian Ocean outbreak enhanced the ability of CHIKV replication in mosquitoes’ vectors (Tsetsarkin *et al.*, 2007).




Figure 2.1 Genome of CHIKV (Niyas *et al.*, 2010).




Figure 2.2 Phylogenetic tree of CHIKV (Lanciotti and Valadere, 2014)

2.3 E2 protein of CHIKV

E2 envelope protein in *alphavirus* which is a type I transmembrane glycoprotein is responsible for the receptor binding (Kielian and Rey, 2006; Brehin *et al.*, 2008). E2 protein consists of three domains A, B and C. Domain C is situated close to the membrane of virus, domain A are positioned in the center of the protein and domain B are at the distal end, which prominently visible on the surface of the virus (Voss *et al.*, 2010; Li *et al.*, 2010). Domain A and B contain most of the residues that affect cell attachment and cell tropism (Voss *et al.*, 2010). Based on the study of Weger-Lucarelli *et al.* (2016), the changed in domains A or C results in attenuated of the viruses, hence this showed that each of the domains of E2 protein have an important role in viral infection. Thus, E2 protein of CHIKV have been a target for recombinant subunit vaccine development and IgM or IgG based diagnostic assay (Verma *et al.*, 2016).

2.4 Epidemiology of CHIKV

CHIKV are probably originated from Central/East Africa, where the virus is circulating in a sylvatic cycle between forest-dwelling *Aedes* species mosquitoes and wild primate such as chimpanzees, monkeys and baboons (Powers *et al.*, 2000). The first major urban outbreak of chikungunya fever was recorded in the early 1960s in Bangkok (Nimmannitya *et al.*, 1969) and from 1963-1971 in India (Nimmannitya *et al.*, 1969; Shah, Gibbs Jr and Banerjee, 1964). After that, only minor outbreaks occurred occasionally over the next 30 years. However, CHIKV re-emerged during 2004 at Kenya and spread throughout countries around the Indian Ocean region, Southeast Asia and major epidemic outbreaks at La Réunion a French island from 2005 to 2006 (Renault *et al.*, 2007). During 2004 to 2007 the outbreaks happened continuously with hundreds of thousands of reported cases and novel geographical areas such as Europe (UK, Belgium, Germany, Czech Republic, Norway, Italy, Spain and France), Hong Kong, Canada, Taiwan, Sri

Lanka and USA (Powers and Logue, 2007). These reported cases in new geographical area (Figure 2.3) were directly associated with the return of tourists from India and island of Indian Ocean (Warner *et al.*, 2006).

The first CHIKV outbreak in Malaysia was documented during 1998-1999 at Port Klang which affected more than 51 persons (Lam *et al.*, 2001) followed by the outbreak in Bagan Panchor, Perak (Kumarasamy *et al.*, 2006), and Kinta district, Perak in 2006 (Noridah *et al.*, 2007). However, the third and largest outbreak was documented in Tangkak, Johor in 2008 (Apandi *et al.*, 2011).

Geographical Distribution of Chikungunya Virus




Figure 2.3 Worldwide distribution of CHIKV (Powers and Logue, 2007).

2.5 Clinical symptoms of CHIKV infection

The incubation period of CHIK fever is generally 2 – 10 days (Burt *et al.*, 2012; Singh and Unni, 2011). Some of the seroprevalence studies carried out and showed different rates of asymptomatic cases that ranges from 3.8% - 27.7% (Thiberville *et al.*, 2013). According to Win *et al.* (2010), during the acute stages of illness, patients with symptomatic cases typically shows the characteristics of high fever, polyarthralgia, headache, back pain and fatigue within 4 – 7 days of infection. The most distinctive symptom is polyarthralgia which is described in 87% - 98% of the symptomatic cases (Thiberville *et al.*, 2013). After the acute stages, some of the patients' experience recur or consistent symptoms such as arthralgia or musculoskeletal pains as a long-lasting sign (Borgherini *et al.*, 2008). Some of the patients infected with CHIKV in Réunion island also presented with chronic rheumatic manifestation (Sissoko *et al.*, 2009).

In children the sign of CHIK fever are specific such as hyperpigmentation, generalized erythema, maculopapular rash, encephalitis seizures and meningeal syndrome (Robin *et al.*, 2008). Besides that, vertical transmission of virus from mother to children are observed at the outbreak on Réunion island with severe illness observed in 53% of newborns that comprises of encephalopathy with persistent disabilities in 44% of newborns (Gerardin *et al.*, 2008).

2.6 Diagnosis of CHIKV infection

Few methods are used to diagnose CHIKV infection which are through the detection of IgM and IgG antibodies through enzyme linked immunosorbent assay (ELISA) and identification through viral isolation and viral RNA identification through real-time PCR (RT-PCR), but these methods are not carried out widely in hospitals because these methods require special equipment and laboratory skills (Okabayashi *et al.*, 2015). The validation of CHIKV isolation may be carried out through immunofluorescence or RT-PCR. RT-PCR can deliver a quick and sensitive

diagnosis, but it is not widely performed in hospitals because it requires specialist equipment and laboratory skills (Okabayashi *et al.*, 2015). However, this method is important for early diagnosis of CHIKV infection especially newborns with meningoencephalitis and vesiculobullous dermatitis (da Cunha and Trinta, 2017). An anti-CHIKV IgM kit is used in evaluation of patient with suspected CHIKV (Mourya and Mishra, 2006), but the IgM detection kit is only limited to the acute stages of illness (days 1 to 5) (Blacksell *et al.*, 2011). Thus, the development of new antigen based diagnostic assay is essential for a fast and reliable clinical diagnosis (Okabayashi *et al.*, 2015).

2.7 Treatment of CHIKV

Currently there are no specific treatment for CHIKV induced disease, and the treatment plans involve only supportive and symptomatic care (Goupil and Mores, 2016). The treatment for acute stages of infection consists of rest analgesics and antipyretics which includes the administration of non-steroidal anti-inflammatory drugs (NSAIDs) and chloroquine in case of refractory arthralgia (Simon *et al.*, 2007; Alla and Combe, 2011). NSAIDs are usually used to treat inflammation but administration of high doses of NSAIDs can control arthralgia but could cause thrombocytopenia, gastrointestinal bleeding, vomiting, gastritis and nausea (Pialoux *et al.*, 2007).

Some small molecule drugs that target the viral replication are still in experimental stages such as ribavirin, chloroquine and Arbidol. Ribavirin shows antiviral activity against a various type of RNA virus especially viruses in the genus *Alphavirus*, such as Semliki forest virus (SFV) and CHIKV (Ho, Ng and Chu, 2010). Currently, ribavirin is used in combination with interferon- α to treat hepatitis C virus (HPC) infection and respiratory syncytial virus (RSV) infections (Crotty, Cameron and Andino, 2002). Next, chloroquine are weak bases that could

affect acid vesicles and could causes the dysfunction of some enzymes. Some of the virus enter their host cell through endocytosis in lysosomal compartment and the low pH disrupts the viral particle and releases the viral nucleic acid. Thus, chloroquine can impair the replication of viruses by interacting with endosome mediated viral entry (Savarino *et al.*, 2003). The application of chloroquine as a treatment for viruses was documented more than 40 years ago (Inglot, 1969). However, for *alphavirus* SFV, the study of Maheshwari, Srikantan and Bhartiya (1991), shows that chloroquine enhanced virus replication in mice. Besides, the antiviral drug Arbidol (ARB) that has developed by the Center for Drug Chemistry, Moscow is effective against respiratory virus such as influenza A and B viruses and respiratory syncytial virus (RSV) (Delogu *et al.*, 2011; Brooks *et al.*, 2012). A study also shows that ARB was effective against CHIKV *in vitro* by interfering with the early stages of infection such as the virus attachment and entry by targeting the cellular membrane (Delogu *et al.*, 2011).

2.8 Serological test

Serological test is effective in diagnosing infectious diseases. Through the principle of reaction of antigen-antibody, it is used to detect serum antibodies or antigen (Kim, Hwang and Oh, 2016). Serology is the detection of increasing titers of antibodies between an acute and convalescent stage of infection or detection of IgM in primary infection. There are few serological techniques that could be used which includes, ELISA, agglutination, precipitation, complement-fixation, fluorescent antibodies or western blot (Martinaskova and Valentova, 2012).

3.0 MATERIALS AND METHODS

3.1 LB medium & agar plate preparation

Luria Bertani (LB) medium consists of 5 g tryptone (BIO 101, CA, USA), 2.5 g yeast extract (amresco, OH, USA), 5 g NaCl (R&M, U.K) are dissolved with R.O water and the volume was adjusted to 500 ml. Then the media were autoclaved at 121 °C, 15 psi for 20 minutes (Hirayama, Japan). After the LB medium has cooled down, 0.5 ml of 50 µg/ml kanamycin (Appendix A.1.3) were added in biosafety cabinet (Envair, U.K) into the medium and stored at 4 °C until used. Then LB agar plate consists of 5 g tryptone (BIO 101, CA, USA), 2.5 g yeast extract (amresco, OH, USA), 5 g NaCl (R&M, U.K), 7.5g agar were dissolved with R. O water and the volume was adjusted to 500 ml. Then the LB agar were autoclaved and 0.5 ml of 50 µg/ml kanamycin (Appendix A.1.3) were added after the agar was cooled down to around 50 °C. After that the LB agar was poured into 20 different plates.

3.2 Bacteria culture growth

The colonies of CHIKV E2 in *E. coli* labelled with BS1/pE2/Mach1/COL1, BS1/pE2/Mach1/COL, BS1/pE2/Mach1/COL5, BS1/pE2/Mach1/COL6 and BS1/pE2/Mach1/COL7 together with the vector DNA colony pET SUMO/Mach1/COL4 were streaked on Luria Bertani (LB) agar plate containing 50 µg/ml kanamycin in a biosafety cabinet (Envair, U.K). The plates were incubated overnight at 37 °C in an incubator (Jisico, Korea).

3.3 Colony polymerase chain reaction (PCR)

First, colony PCR was done to check the presence of the E2 protein in the colonies. An individual colony was picked from plates by using a sterile toothpick and mixed into a 50 µl PCR reagent mixture that contain (5 µl 10X Taq buffer, 3 µl 25 mM MgCl₂, 1.5 µl pE2_F1 primer (20 pmol/µl), 1.5 µl pET SUMO reverse primer (10 pmol/µl), 1 µl 10 mM dNTP

(Fermentas, NY, USA), 0.5 µl Taq polymerase (Fermentas, NY, USA) and 37.5 µl UHQ water). The primer sequences used were shown in Table 3.1. For the pET SUMO control, the colony was picked and mixed in a 50 µl PCR reagent mixture that contain (5 µl 10X Taq buffer, 3 µl 25 mM MgCl₂, 1.5 µl pET SUMO forward primer (10 pmol/µl), 1.5 µl pET SUMO reverse primer (10 pmol/µl), 1 µl 10mM dNTP (Fermentas, NY, USA), 0.5 µl Taq polymerase (Fermentas, NY, USA), 37.5 µl UHQ water). The PCR was then performed in a thermocycler (PTC-200, MJ Research, NY, USA) in the condition shown in Table 3.2. Then, 1.5% agarose gel electrophoresis was run at 120 V for 40 minutes (Thermo Scientific, NH, USA) and the image was captured (Bio-Rad, CA, USA).

Table 3.1 Primers for colony PCR

Primers	Primer sequences
pET SUMO Forward	5'- AGA TTC TTG TAC GAC GGT ATT AG – 3'
pET SUMO Reverse	5'- TAG TTA TTG CTC AGC GGT GG – 3'

Table 3.2 Condition for colony PCR reaction

Condition	Temperature (°C)	Duration	Cycles
Initial denaturation	94	8 minutes	1
Denaturation	94	20 seconds	35
Annealing	55	30 seconds	
Extension	72	2 minutes	
Final extension	72	5 minutes	1